


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.


In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



Contents lists available at ScienceDirect

Insurance: Mathematics and Economics

journal homepage: www.elsevier.com/locate/ime

On multivariate extensions of Conditional-Tail-Expectation

Areski Cousin^{a,1}, Elena Di Bernardino^{b,*}^a Université de Lyon, Université Lyon 1, ISFA, Laboratoire SAF, 50 avenue Tony Garnier, 69366 Lyon, France^b CNAM, Paris, Département IMATH, Laboratoire Cédric EA4629, 292 rue Saint-Martin, Paris Cedex 03, France

ARTICLE INFO

Article history:

Received November 2013

Received in revised form

January 2014

Accepted 25 January 2014

Keywords:

Multivariate risk measures

Level sets of distribution functions

Multivariate probability integral transformation

Stochastic orders

Copulas and dependence

ABSTRACT

In this paper, we introduce two alternative extensions of the classical univariate *Conditional-Tail-Expectation* (CTE) in a multivariate setting. The two proposed multivariate CTEs are vector-valued measures with the same dimension as the underlying risk portfolio. As for the multivariate Value-at-Risk measures introduced by Cousin and Di Bernardino (2013), the *lower-orthant CTE* (resp. the *upper-orthant CTE*) is constructed from level sets of multivariate distribution functions (resp. of multivariate survival distribution functions). Contrary to allocation measures or systemic risk measures, these measures are also suitable for multivariate risk problems where risks are heterogeneous in nature and cannot be aggregated together. Several properties have been derived. In particular, we show that the proposed multivariate CTEs satisfy natural extensions of the positive homogeneity property, the translation invariance property and the comonotonic additivity property. Comparison between univariate risk measures and components of multivariate CTE is provided. We also analyze how these measures are impacted by a change in marginal distributions, by a change in dependence structure and by a change in risk level. Sub-additivity of the proposed multivariate CTE-s is provided under the assumption that all components of the random vectors are independent. Illustrations are given in the class of Archimedean copulas.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

As illustrated by the recent financial turmoil, risks are strongly interconnected. Consequently, risk quantification in multivariate settings has recently been the subject of great interest. Much research has been devoted to construction of risk measures that account both for marginal effects and dependence between risks.

In the literature, several generalizations of the classical univariate Conditional-Tail-Expectation (CTE) have been proposed, mainly using as conditioning events the total risk or some extreme risks. These measures can be used as capital allocation rules for financial institutions. The aim is to find the contribution of each subsidiary (or risk category) to the total economic capital. As can be seen in Scaillet (2004) and Tasche (2008), the Euler or Shapley–Aumann allocation rule associated with a particular univariate risk measure (such as VaR or CTE) involves the dependence structure between marginal and aggregated risks. More formally, let $\mathbf{X} =$

(X_1, \dots, X_d) represent the risk exposures of a given financial institution, where, for any $i = 1, \dots, d$, the component X_i denotes the marginal risk (usually claim or loss) associated with the underlying entity i (the latter could be, for instance, a subsidiary, an operational branch or a risk category). Then, the sum $S = X_1 + \dots + X_d$ corresponds to the company aggregated risk, whereas $X_{(1)} = \min\{X_1, \dots, X_d\}$ and $X_{(d)} = \max\{X_1, \dots, X_d\}$ are the extreme risks. In capital allocation problems, we are not only interested in the “stand-alone” risk measures $\text{CTE}_\alpha(X_i) = \mathbb{E}[X_i | X_i > Q_{X_i}(\alpha)]$, where $Q_{X_i}(\alpha) = \inf\{x \in \mathbb{R}_+ : F_{X_i}(x) \geq \alpha\}$ is the univariate quantile function of X_i at risk level α , but also in

$$\text{CTE}_\alpha^{\text{sum}}(X_i) = \mathbb{E}[X_i | S > Q_S(\alpha)], \quad (1)$$

$$\text{CTE}_\alpha^{\text{min}}(X_i) = \mathbb{E}[X_i | X_{(1)} > Q_{X_{(1)}}(\alpha)], \quad (2)$$

$$\text{CTE}_\alpha^{\text{max}}(X_i) = \mathbb{E}[X_i | X_{(d)} > Q_{X_{(d)}}(\alpha)], \quad (3)$$

for $i = 1, \dots, d$. The interested reader is referred to Cai and Li (2005) for further details. An explicit formula for $\text{CTE}_\alpha^{\text{sum}}(X_i)$ is provided in Landsman and Valdez (2003) in the case of elliptical distribution functions, Cai and Li (2005) for *phase-type* distributions and in Bargès et al. (2009) for Farlie–Gumbel–Morgenstern family of copulas. Furthermore, we recall that $\text{CTE}_\alpha^{\text{sum}}(X_i)$ corresponds to the “Euler allocation rule” associated with the univariate CTE (see, e.g., Tasche, 2008).

* Corresponding author. Tel.: +33 647158175.

E-mail addresses: areski.cousin@univ-lyon1.fr (A. Cousin), elena.di_bernardino@cnam.fr (E. Di Bernardino).URLs: <http://www.acousin.net/> (A. Cousin),<http://isfaserveur.univ-lyon1.fr/~elena.dibernardino/> (E. Di Bernardino).¹ Tel.: +33 437287439.

Another problem which recently receives a great interest is the construction of systemic risk measures. One of the proposed measure is the Marginal Expected Shortfall (MES) defined as the expected loss on its equity return (X) conditional on the occurrence of a loss in the aggregated return of the financial sector (Y), i.e.,

$$\text{MES}_\alpha(X) = \mathbb{E}[X \mid Y > Q_Y(\alpha)], \quad (4)$$

where $Q_Y(\alpha)$ is the (α) -th quantile of the distribution of Y . The MES of a financial institution aims at detecting which firms in the economy are the more vulnerable in case of a global financial distress. On mathematical grounds, this measure is similar to the allocation measure CTE^{sum} . The interested reader is referred to Acharya et al. (2010) or Brownlees and Engle (2012) for more details. Cai et al. (2013) propose a non-parametric estimator of the MES using extreme value theory. The CoVaR (conditional VaR) of company i is instead given by

$$\text{CoVaR}_\alpha^i(X) = \text{VaR}_\alpha(S \mid X \geq \text{VaR}_\alpha(X)). \quad (5)$$

As opposed to the MES, the CoVaR (conditional VaR) is constructed in order to identify which firms in the economy have a great importance in terms of systemic risk (see Adrian and Brunnermeier, 2011 or Mainik and Schaanning, 2012).

However, the previous risk measures are not suitable for multivariate risk problem where risks are heterogeneous in nature and thus cannot be aggregated together or even compared. This is the case for instance for risks which are difficult to expressed under the same numéraire or when one has to deal with non-monetary risks or exogenous risks. The literature which deals with risk measures for intrinsically multivariate problems can be divided in two categories. The first group of papers are interested in extending classical univariate axioms to different multivariate settings (see for instance Jouini et al., 2004, Burgert and Rüschendorf, 2006, Rüschendorf, 2006, Cascos and Molchanov, 2007, Hamel and Heyde, 2010 and Ekeland et al., 2012). One of the objectives is to derive theoretical representation of risk measures. This is done without proposing tractable constructions for the axiom-consistent multivariate measures. Another group of papers investigates different generalizations of the concept of quantiles in a multivariate setting. Unsurprisingly, the main difficulty regarding multivariate generalizations of quantile-based risk measures (as the VaR and the CTE) is the fact that vector preorders are, in general, partial preorders. Then, what can be considered in a context of multidimensional portfolios as the analogous of a “worst case” scenario and a related “tail distribution”? For example, Massé and Theodorescu (1994) define multivariate α -quantiles for bivariate distribution as the intersection of half-planes whose the distribution is at least equal to α . Koltchinskii (1997) provides a general treatment of multivariate quantiles as inversions of mappings. Another approach is to use geometric quantiles (see, for example, Chaouch et al., 2009). Along with the geometric quantile, the notion of depth function has been developed in recent years to characterize the quantile of multidimensional distribution functions (for further details see, for instance, Chauvigny et al., 2011). We refer to Serfling (2002) for a review of multivariate quantiles.

When it turns to generalize the Value-at-Risk measure, Embrechts and Puccetti (2006), Nappo and Spizzichino (2009), Prékopa (2012) use the notion of quantile curve but these papers do not investigate whether these measures are compatible with some desirable axioms. Moreover, the proposed risk measures are hyperspaces and thus quantify a vector of risks with an infinite number of points. Contrarily to the latter approach, the multivariate Conditional-Tail-Expectation proposed in this paper quantifies multivariate risks in a more parsimonious and synthetic way. This feature can be relevant for operational applications since it can ease discrimination between portfolio of risks. Lee and Prékopa (2013) introduce a real-valued measure of multivariate risks which also bears on quantile curves but the proposed measure relies on a somehow arbitrary convex combination.

We propose two vector-valued extensions of the univariate Conditional-Tail-Expectation. The lower-orthant CTE of a random vector X (introduced by Di Bernardino et al., 2013 in a bivariate setting) is defined as the conditional expectation of X given that the latter is located in the α -upper level set of its distribution function. The upper-orthant CTE of X is defined as the conditional expectation of X given that the latter is in the $(1 - \alpha)$ -lower level set of its survival function. Several properties have been derived. We provide an integral representation of the proposed measures in terms of the multivariate VaR introduced in Cousin and Di Bernardino (2013) and we show that the proposed multivariate CTE-s satisfy natural extensions of the positive homogeneity property, the translation invariance property and the comonotonic additivity property. We show that the proposed measures are sub-additive for independent vectors with independent components. We also provide comparisons between univariate risk measures and components of the proposed multivariate CTE. We analyze how these measures are impacted by a change in marginal distributions, by a change in dependence structure and by a change in risk level.

The paper is organized as follows. In Section 2, we give the definition of the lower-orthant and the upper-orthant Conditional-Tail-Expectation measures. We then show that these measures satisfy multivariate extensions of Artzner et al. (1999)’s invariance properties (see Section 2.1). Illustrations in some Archimedean copula cases are presented in Section 2.2. We also compare the components of these multivariate CTE measures with the associated univariate VaR, the associated univariate CTE and with the multivariate lower-orthant and upper-orthant VaR previously introduced by Cousin and Di Bernardino (2013) (see Section 2.3). The behavior of our CTE-s with respect to a change in marginal distributions, a change in dependence structure and a change in risk level α is discussed respectively in Sections 2.4–2.6. The conclusion discusses open problems and possible directions for future work.

2. Multivariate generalization of the Conditional-Tail-Expectation measure

As in the univariate case, the multivariate VaR introduced in Cousin and Di Bernardino (2013) does not give any information regarding the upper tail of the loss distribution function and especially its degree of thickness above the VaR threshold. In an univariate setting, the problem has been overcome by considering for instance the Conditional-Tail-Expectation (CTE) risk measure,² which is defined as the conditional expectation of losses given that the latter exceed VaR. Following Artzner et al. (1999), the CTE at level α for a distribution function F (or survival function \bar{F}) is given by

$$\text{CTE}_\alpha(X) := \mathbb{E}[X \mid X \geq \text{VaR}_\alpha(X)], \quad (6)$$

where $\text{VaR}_\alpha(X)$ is the univariate Value-at-Risk defined by

$$\begin{aligned} \text{VaR}_\alpha(X) &:= \inf \{x \in \mathbb{R} : F(x) \geq \alpha\} \\ &= \inf \{x \in \mathbb{R} : \bar{F}(x) \leq 1 - \alpha\}. \end{aligned}$$

Since the sets $\{X \geq \text{VaR}_\alpha(X)\}$, $\{F(X) \geq \alpha\}$ and $\{\bar{F}(X) \leq 1 - \alpha\}$ correspond to the same event in a univariate setting, the CTE can alternatively be defined³ as

$$\text{CTE}_\alpha(X) := \mathbb{E}[X \mid F(X) \geq \alpha] = \mathbb{E}[X \mid \bar{F}(X) \leq 1 - \alpha]. \quad (7)$$

The CTE can then be viewed as the conditional expectation of X given that X falls into the α -lower-level set of its distribution function $\underline{L}(\alpha) := \{x \in \mathbb{R}_+ : F(x) \geq \alpha\}$ or equivalently in the $(1 - \alpha)$ -upper-level set of its survival function $\bar{L}(\alpha) := \{x \in \mathbb{R}_+ : \bar{F}(x) \leq$

² This measure is also called Tail Conditional Expectation. As far as continuous distribution functions are considered, the CTE measure is coherent in the sense of Artzner’s axioms and it coincides with the Expected Shortfall or Tail VaR.

³ Note that this definition does not depend on VaR.

$1 - \alpha$ }. This definition can naturally be extended in higher dimension, keeping in mind that the two previous sets $\underline{L}(\alpha)$ and $\bar{L}(\alpha)$ are different in general as soon as the dimension d of the underlying risk vector is greater or equal to 2. In analogy with Embrechts and Puccetti (2006) notations, we will denote by $\underline{\text{CTE}}$ the lower-orthant Conditional-Tail-Expectation based on level sets of the underlying distribution function (distribution function measures mass of probabilities in lower-orthant regions) and by $\overline{\text{CTE}}$ the upper-orthant Conditional-Tail-Expectation based on level sets of the underlying survival function (survival function measures mass of probabilities in upper-orthant regions).

Assumption 2.1. In the following, we will consider non-negative absolutely-continuous random vector⁴ $\mathbf{X} = (X_1, \dots, X_d)$ (with respect to Lebesgue measure λ on \mathbb{R}^d) with partially increasing multivariate distribution function⁵ F and such that $\mathbb{E}[X_i] < \infty$, for $i = 1, \dots, d$. These conditions will be called *regularity conditions*.

Definition 2.1 (Multivariate Lower-Orthant and Upper-Orthant Conditional-Tail-Expectation). Consider a random vector $\mathbf{X} = (X_1, \dots, X_d)$ with distribution function F and survival function \bar{F} , such that $\mathbb{E}[X_i]$ is finite for all $i = 1, \dots, d$. For $\alpha \in (0, 1)$, we define the lower-orthant Conditional-Tail-Expectation at probability level α by

$$\underline{\text{CTE}}_\alpha(\mathbf{X}) = \mathbb{E}[\mathbf{X} \mid F(\mathbf{X}) \geq \alpha] = \begin{pmatrix} \mathbb{E}[X_1 \mid F(\mathbf{X}) \geq \alpha] \\ \vdots \\ \mathbb{E}[X_d \mid F(\mathbf{X}) \geq \alpha] \end{pmatrix},$$

and the upper-orthant Conditional-Tail-Expectation at probability level α by

$$\overline{\text{CTE}}_\alpha(\mathbf{X}) = \mathbb{E}[\mathbf{X} \mid \bar{F}(\mathbf{X}) \leq 1 - \alpha] = \begin{pmatrix} \mathbb{E}[X_1 \mid \bar{F}(\mathbf{X}) \leq 1 - \alpha] \\ \vdots \\ \mathbb{E}[X_d \mid \bar{F}(\mathbf{X}) \leq 1 - \alpha] \end{pmatrix}.$$

Remark that the lower-orthant CTE is a multivariate generalization of the bivariate Conditional-Tail-Expectation previously introduced in Di Bernardino et al. (2013).

From now on, we denote by $\underline{\text{CTE}}_\alpha^1(\mathbf{X}), \dots, \underline{\text{CTE}}_\alpha^d(\mathbf{X})$ the components of the vector $\underline{\text{CTE}}_\alpha(\mathbf{X})$ and by $\overline{\text{CTE}}_\alpha^1(\mathbf{X}), \dots, \overline{\text{CTE}}_\alpha^d(\mathbf{X})$ the components of the vector $\overline{\text{CTE}}_\alpha(\mathbf{X})$.

Note that if \mathbf{X} is an exchangeable random vector, $\underline{\text{CTE}}_\alpha^i(\mathbf{X}) = \underline{\text{CTE}}_\alpha^j(\mathbf{X})$ and $\overline{\text{CTE}}_\alpha^i(\mathbf{X}) = \overline{\text{CTE}}_\alpha^j(\mathbf{X})$ for any $i, j = 1, \dots, d$. Furthermore, given a univariate random variable X , $\mathbb{E}[X \mid F_X(X) \geq \alpha] = \mathbb{E}[X \mid F_X(X) \leq 1 - \alpha] = \text{CTE}_\alpha(X)$, for all α in $(0, 1)$. Hence, lower-orthant and upper-orthant Conditional-Tail-Expectation coincide with the usual CTE in the particular case where $d = 1$.

Let us remark that, under the regularity assumption, the multivariate lower-orthant (resp. upper-orthant) Conditional-Tail-Expectation can be represented as an integral transformation of the multivariate lower-orthant VaR (resp. upper-orthant VaR) introduced by Cousin and Di Bernardino (2013), i.e.,

$$\underline{\text{CTE}}_\alpha^i(\mathbf{X}) = \frac{1}{1 - K(\alpha)} \int_\alpha^1 \text{VaR}_\gamma^i(\mathbf{X}) K'(\gamma) d\gamma, \quad (8)$$

⁴ We restrict ourselves to \mathbb{R}_+^d because, in our applications, components of d -dimensional vectors correspond to random losses and are then valued in \mathbb{R}_+ .

⁵ A function $F(x_1, \dots, x_d)$ is partially increasing on $\mathbb{R}_+^d \setminus (0, \dots, 0)$ if the functions of one variable $g(\cdot) = F(x_1, \dots, x_{j-1}, \cdot, x_{j+1}, \dots, x_d)$ are increasing. About properties of partially increasing multivariate distribution functions we refer the interested reader to Rossi (1973) and Tibiletti (1991).

and

$$\overline{\text{CTE}}_\alpha^i(\mathbf{X}) = \frac{1}{\widehat{K}(1 - \alpha)} \int_\alpha^1 \text{VaR}_\gamma^i(\mathbf{X}) \widehat{K}'(1 - \gamma) d\gamma, \quad (9)$$

where K is the Kendall distribution of X , i.e., $K(x) = \mathbb{P}(F(\mathbf{X}) \leq x)$, for all x in $(0, 1)$ and \widehat{K} is its “upper-orthant” Kendall distribution, i.e., $\widehat{K}(x) = \mathbb{P}(\bar{F}(\mathbf{X}) \leq x)$, for all x in $(0, 1)$. Formula (8) and (9) will be useful in Proposition 2.8 and Corollary 2.3 below.

Remark that the existence of the density K' and \widehat{K}' that appears in Eqs. (8)–(9) is guaranteed by the *regularity conditions* (for further details, see Proposition 1 in Imlahi et al., 1999 or Proposition 4 in Chakak and Ezzer, 2000). The interested reader is also referred to Cousin and Di Bernardino (2013).

2.1. Invariance properties

In the present section, the aim is to analyze the lower-orthant and the upper-orthant CTE introduced in Definition 2.1 in terms of classical invariance properties of risk measures (we refer the interested reader to Artzner et al., 1999). In analogy with Section 2.1 in Cousin and Di Bernardino (2013), we now introduce the following results (Proposition 2.1 and Corollary 2.1) that will be useful in order to prove invariance properties of our risk measures.

Proposition 2.1. Let the function h be such that $h(x_1, \dots, x_d) = (h_1(x_1), \dots, h_d(x_d))$.

– If h_1, \dots, h_d are non-decreasing functions, then the following relations hold

$$\underline{\text{CTE}}_\alpha^i(h(\mathbf{X})) = \mathbb{E}[h_i(X_i) \mid F_{\mathbf{X}}(\mathbf{X}) \geq \alpha], \quad i = 1, \dots, d$$

and

$$\overline{\text{CTE}}_\alpha^i(h(\mathbf{X})) = \mathbb{E}[h_i(X_i) \mid \bar{F}_{\mathbf{X}}(\mathbf{X}) \leq 1 - \alpha], \quad i = 1, \dots, d.$$

– If h_1, \dots, h_d are non-increasing functions, then the following relations hold

$$\underline{\text{CTE}}_\alpha^i(h(\mathbf{X})) = \mathbb{E}[h_i(X_i) \mid \bar{F}_{\mathbf{X}}(\mathbf{X}) \geq \alpha], \quad i = 1, \dots, d$$

and

$$\overline{\text{CTE}}_\alpha^i(h(\mathbf{X})) = \mathbb{E}[h_i(X_i) \mid F_{\mathbf{X}}(\mathbf{X}) \leq 1 - \alpha], \quad i = 1, \dots, d.$$

Proof. From Definition 2.1, $\underline{\text{CTE}}_\alpha^i(h(\mathbf{X})) = \mathbb{E}[h_i(X_i) \mid F_{h(\mathbf{X})}(h(\mathbf{X})) \geq \alpha]$, for $i = 1, \dots, d$. Since

$$F_{h(\mathbf{X})}(y_1, \dots, y_d) = \begin{cases} F_{\mathbf{X}}(h^{-1}(y_1), \dots, h^{-1}(y_d)), & \text{if } h_1, \dots, h_d \text{ are non-decreasing} \\ & \text{functions,} \\ \bar{F}_{\mathbf{X}}(h^{-1}(y_1), \dots, h^{-1}(y_d)), & \text{if } h_1, \dots, h_d \text{ are non-increasing} \\ & \text{functions,} \end{cases}$$

then we obtain the result. \square

Finally, we can state the following result that proves positive homogeneity and translation invariance for our measures.

Proposition 2.2. For $\alpha \in (0, 1)$, the multivariate upper-orthant and lower-orthant Conditional-Tail-Expectation satisfy the following properties:

Positive Homogeneity: $\forall \mathbf{c} = (c_1, \dots, c_d)' \in \mathbb{R}_+^d$,

$$\underline{\text{CTE}}_\alpha(c_1 X_1, \dots, c_d X_d) = (c_1 \underline{\text{CTE}}_\alpha^1(\mathbf{X}), \dots, c_d \underline{\text{CTE}}_\alpha^d(\mathbf{X}))'$$

$$\overline{\text{CTE}}_\alpha(c_1 X_1, \dots, c_d X_d) = (c_1 \overline{\text{CTE}}_\alpha^1(\mathbf{X}), \dots, c_d \overline{\text{CTE}}_\alpha^d(\mathbf{X}))'$$

Translation Invariance: $\forall \mathbf{c} \in \mathbb{R}_+^d$,

$$\underline{\text{CTE}}_\alpha(\mathbf{c} + \mathbf{X}) = \mathbf{c} + \underline{\text{CTE}}_\alpha(\mathbf{X}), \quad \overline{\text{CTE}}_\alpha(\mathbf{c} + \mathbf{X}) = \mathbf{c} + \overline{\text{CTE}}_\alpha(\mathbf{X}).$$

The proof comes down from Proposition 2.1.

Remark 1. (i) In the univariate as well in the multivariate setting, a risk measure may be interpreted as the riskiness of a portfolio or the amount of capital that should be added to a portfolio with a given loss, so that the portfolio can then be deemed *acceptable*. In this respect, the translation invariance property for (multivariate) risk measures are necessary if the risk-capital interpretation we stated above is to make sense. The homogeneity property is often motivated by a change of currency argument: the amount of required capital in order to manager risks should be independent of the currency in which it is expressed (e.g. see Artzner et al., 1999). For these reasons, the invariance properties proved in Proposition 2.2 play a central role in practical applications.

(ii) It is straightforward to show that the capital allocation measures presented in (1)–(3) do not satisfy the invariance properties stated in Proposition 2.2 as soon as the components of $\mathbf{c} = (c_1, \dots, c_d)$ are all different. In that sense, these measures may not be suitable for intrinsically multivariate problem where risks cannot be expressed under the same numéraire.

Remark 2. For $\alpha = 0$, using Definition 2.1, we obtain

$$\underline{\text{CTE}}_0(\mathbf{X}) = \overline{\text{CTE}}_0(\mathbf{X}) = \begin{pmatrix} \mathbb{E}[X_1] \\ \vdots \\ \mathbb{E}[X_d] \end{pmatrix} = \mathbb{E}[\mathbf{X}].$$

Then, as in the univariate case, the multivariate lower-orthant and upper-orthant CTE-s are equal to the expected value of the underlying random vector for $\alpha = 0$.

Since these two new measures are not the same in general for dimension greater or equal to 2, we also provide some connections between $\underline{\text{CTE}}$ and $\overline{\text{CTE}}$. From Proposition 2.1 one can obtain the following property which links the multivariate upper-orthant Conditional-Tail-Expectation and the lower-orthant one.

Corollary 2.1. Let h be a linear function such that $h(x_1, \dots, x_d) = (h_1(x_1), \dots, h_d(x_d))$.

– If h_1, \dots, h_d are non-decreasing functions then it holds that

$$\underline{\text{CTE}}_\alpha(h(\mathbf{X})) = h(\underline{\text{CTE}}_\alpha(\mathbf{X})) \quad \text{and} \\ \overline{\text{CTE}}_\alpha(h(\mathbf{X})) = h(\overline{\text{CTE}}_\alpha(\mathbf{X})).$$

– If h_1, \dots, h_d are non-increasing functions then it holds that

$$\underline{\text{CTE}}_\alpha(h(\mathbf{X})) = h\left(\frac{\mathbb{E}[\mathbf{X}]}{1 - \widehat{K}(\alpha)} - \frac{\widehat{K}(\alpha)}{1 - \widehat{K}(\alpha)} \overline{\text{CTE}}_{1-\alpha}(\mathbf{X})\right)$$

and

$$\overline{\text{CTE}}_\alpha(h(\mathbf{X})) = h\left(\frac{\mathbb{E}[\mathbf{X}]}{K(1-\alpha)} - \frac{1 - K(1-\alpha)}{K(1-\alpha)} \underline{\text{CTE}}_{1-\alpha}(\mathbf{X})\right),$$

where $K(x) = \mathbb{P}(F(\mathbf{X}) \leq x)$ and $\widehat{K}(x) = \mathbb{P}(\overline{F}(\mathbf{X}) \leq x)$, for all $x \in (0, 1)$.

Example 1. If $X = (X_1, \dots, X_d)$ is a random vector with uniform margins and if, for all $i = 1, \dots, d$, we consider the functions h_i such that $h_i(x) = 1 - x$, $x \in [0, 1]$, then from Corollary 2.1, we get

$$\overline{\text{CTE}}_\alpha^i(\mathbf{X}) = \frac{1}{\widehat{K}(1-\alpha)} \left[\frac{1}{2} - (1 - \widehat{K}(1-\alpha)) \right. \\ \left. \times (1 - \underline{\text{CTE}}_{1-\alpha}^i(\mathbf{1} - \mathbf{X})) \right] \quad (10)$$

for all $i = 1, \dots, d$, where $\mathbf{1} - \mathbf{X} = (1 - X_1, \dots, 1 - X_d)$. Note that \widehat{K} is the Kendall distribution function associated with the vector $\mathbf{1} - \mathbf{X}$.

Additionally, if \mathbf{X} and $\mathbf{1} - \mathbf{X}$ have the same distribution function, then X is invariant in law by central symmetry and the following relation holds:

$$\overline{\text{CTE}}_\alpha^i(\mathbf{X}) = \frac{1}{K(1-\alpha)} \left[\frac{1}{2} - (1 - K(1-\alpha)) \right. \\ \left. \times (1 - \underline{\text{CTE}}_{1-\alpha}^i(\mathbf{X})) \right]. \quad (11)$$

For instance, this property is satisfied for vector \mathbf{X} which follows an elliptical or a Frank copula structure.

We now show that, as in the univariate setting, the lower-orthant and the upper-orthant CTE-s are additive for π -comonotonic couple of random vectors as defined in Puccetti and Scarsini (2010).

Definition 2.2. A couple (\mathbf{X}, \mathbf{Y}) of d -dimensional random vectors is said to be π -comonotonic if there exists a d -dimensional random vector $\mathbf{Z} = (Z_1, \dots, Z_d)$ and non-decreasing functions $f_1, \dots, f_d, g_1, \dots, g_d$ such that

$$(\mathbf{X}, \mathbf{Y}) \stackrel{d}{=} ((f_1(Z_1), \dots, f_d(Z_d)), (g_1(Z_1), \dots, g_d(Z_d))).$$

Proposition 2.3. Let (\mathbf{X}, \mathbf{Y}) be a π -comonotonic couple of random vectors, then

$$\underline{\text{CTE}}_\alpha(\mathbf{X} + \mathbf{Y}) = \underline{\text{CTE}}_\alpha(\mathbf{X}) + \underline{\text{CTE}}_\alpha(\mathbf{Y}), \\ \overline{\text{CTE}}_\alpha(\mathbf{X} + \mathbf{Y}) = \overline{\text{CTE}}_\alpha(\mathbf{X}) + \overline{\text{CTE}}_\alpha(\mathbf{Y}).$$

Proof. We focus on the lower-orthant CTE. Similar arguments apply for the upper-orthant CTE. Let $\mathbf{X} = (X_1, \dots, X_d)$ and $\mathbf{Y} = (Y_1, \dots, Y_d)$ be two π -comonotonic random vectors. There exists a random vector $\mathbf{Z} = (Z_1, \dots, Z_d)$ such that, for any $i = 1, \dots, d$, $X_i = f_i(Z_i)$ and $Y_i = g_i(Z_i)$ where f_i and g_i are non-decreasing functions. Let f be the function defined by $f(x_1, \dots, x_d) = (f_1(x_1), \dots, f_d(x_d))$, g be the function defined by $g(x_1, \dots, x_d) = (g_1(x_1), \dots, g_d(x_d))$ and h be the function defined by $h(x_1, \dots, x_d) = (h_1(x_1), \dots, h_d(x_d))$ where $h_i := f_i + g_i$, $i = 1, \dots, d$. The function h_i , $i = 1, \dots, d$ are non-decreasing as a sum of non-decreasing functions and $\mathbf{X} + \mathbf{Y} = h(\mathbf{Z})$. We obtain from Proposition 2.1 that, for any $i = 1, \dots, d$, $\underline{\text{CTE}}_\alpha^i(\mathbf{X} + \mathbf{Y}) = \mathbb{E}[h_i(Z_i) \mid F_{\mathbf{Z}}(\mathbf{Z}) \geq \alpha] = \mathbb{E}[f_i(Z_i) \mid F_{\mathbf{Z}}(\mathbf{Z}) \geq \alpha] + \mathbb{E}[g_i(Z_i) \mid F_{\mathbf{Z}}(\mathbf{Z}) \geq \alpha]$ where $F_{\mathbf{Z}}$ denotes the distribution function of \mathbf{Z} . Eventually, $\mathbb{E}[f_i(Z_i) \mid F_{\mathbf{Z}}(\mathbf{Z}) \geq \alpha] = \mathbb{E}[f_i(Z_i) \mid F_{f(\mathbf{Z})}(\mathbf{f}(\mathbf{Z})) \geq \alpha] = \underline{\text{CTE}}_\alpha^i(\mathbf{X})$ and $\mathbb{E}[g_i(Z_i) \mid F_{\mathbf{Z}}(\mathbf{Z}) \geq \alpha] = \mathbb{E}[g_i(Z_i) \mid F_{g(\mathbf{Z})}(\mathbf{g}(\mathbf{Z})) \geq \alpha] = \underline{\text{CTE}}_\alpha^i(\mathbf{Y})$, which concludes the proof. \square

Remark 3. The previous proposition also holds for the lower-orthant VaR and the upper-orthant VaR introduced in Cousin and Di Bernardino (2013).

We now prove that the proposed multivariate measures are subadditive for independent random vectors with independent components.

Proposition 2.4. Let \mathbf{X} and \mathbf{Y} be two d -dimensional independent random vectors with finite expectations for each marginal. If all components in \mathbf{X} and \mathbf{Y} are independent, then the lower-orthant Conditional-Tail-Expectation $\underline{\text{CTE}}$ is such that

$$\underline{\text{CTE}}_\alpha(\mathbf{X} + \mathbf{Y}) \leq \underline{\text{CTE}}_\alpha(\mathbf{X}) + \underline{\text{CTE}}_\alpha(\mathbf{Y}), \quad \text{for } \alpha \in (0, 1).$$

The same result holds for the lower-orthant Conditional-Tail-Expectation $\overline{\text{CTE}}$.

Proof. It is equivalent to prove that, for all $i = 1, \dots, d$ and for all $\alpha \in (0, 1)$,

$$\mathbb{E}[X_i + Y_i \mid F_{\mathbf{X}+\mathbf{Y}}(\mathbf{X} + \mathbf{Y}) \geq \alpha] \leq \mathbb{E}[X_i \mid F_{\mathbf{X}}(\mathbf{X}) \geq \alpha] \\ + \mathbb{E}[Y_i \mid F_{\mathbf{Y}}(\mathbf{Y}) \geq \alpha]. \quad (12)$$

Eq. (12) can be written as:

$$\frac{\mathbb{E}[X_i + Y_i, F_{\mathbf{X}+\mathbf{Y}}(\mathbf{X} + \mathbf{Y}) \geq \alpha]}{\mathbb{P}[F_{\mathbf{X}+\mathbf{Y}}(\mathbf{X} + \mathbf{Y}) \geq \alpha]} \leq \frac{\mathbb{E}[X_i, F_{\mathbf{X}}(\mathbf{X}) \geq \alpha]}{\mathbb{P}[F_{\mathbf{X}}(\mathbf{X}) \geq \alpha]} + \frac{\mathbb{E}[Y_i, F_{\mathbf{Y}}(\mathbf{Y}) \geq \alpha]}{\mathbb{P}[F_{\mathbf{Y}}(\mathbf{Y}) \geq \alpha]}.$$

Note that $X_i + Y_i$ and $X_j + Y_j$ are independent for all $i, j = 1, \dots, d$, with $i \neq j$. Then all denominators of the terms above are the same, i.e., the survival Kendall distribution $\bar{K}(\alpha)$ associated to the d -dimensional independence copula structure. As a result, we have then to prove that, for all $i = 1, \dots, d$ and for all $\alpha \in (0, 1)$,

$$\mathbb{E}[X_i + Y_i, F_{\mathbf{X}+\mathbf{Y}}(\mathbf{X} + \mathbf{Y}) \geq \alpha] \leq \mathbb{E}[X_i, F_{\mathbf{X}}(\mathbf{X}) \geq \alpha] + \mathbb{E}[Y_i, F_{\mathbf{Y}}(\mathbf{Y}) \geq \alpha].$$

Let us denote by $\mathbf{U} = (U_1, \dots, U_d)$ the random vector with components $U_j := F_{X_j+Y_j}(X_j + Y_j)$, $j = 1, \dots, d$ and by $\mathbf{V} = (V_1, \dots, V_d)$ the random vector with components $V_j := F_{X_j}(X_j)$, $j = 1, \dots, d$. As the components of $\mathbf{X} + \mathbf{Y}$ are independent, we get:

$$\mathbb{E}[X_i + Y_i, F_{\mathbf{X}+\mathbf{Y}}(\mathbf{X} + \mathbf{Y}) \geq \alpha] = \mathbb{E}[X_i + Y_i, F_{X_i+Y_i}(X_i + Y_i) \geq \tilde{\alpha}_i]$$

where

$$\tilde{\alpha}_i := \frac{\alpha}{\prod_{j=1, j \neq i}^d U_j}.$$

Using the sub-additivity of the univariate Conditional-Tail-Expectation (see for instance Denuit et al., 2005), the following relation holds

$$\mathbb{E}[X_i + Y_i, F_{X_i+Y_i}(X_i + Y_i) \geq \tilde{\alpha}_i | U_{j, j \neq i}] \leq \mathbb{E}[X_i, F_{X_i}(X_i) \geq \tilde{\alpha}_i | U_{j, j \neq i}] + \mathbb{E}[Y_i, F_{Y_i}(Y_i) \geq \tilde{\alpha}_i | U_{j, j \neq i}]. \quad (13)$$

Note that the vector $(U_1, \dots, X_i, \dots, U_d)$ has the same distribution as the vector $(V_1, \dots, X_i, \dots, V_d)$ since they both share the same marginals and the same dependence structure (the independent copula). As a result, $\mathbb{E}[X_i, F_{X_i}(X_i) \geq \tilde{\alpha}_i | U_{j, j \neq i}] = \mathbb{E}[X_i, F_{\mathbf{X}}(\mathbf{X}) \geq \alpha | V_{j, j \neq i}]$. Using the same argument for the second term in the right hand side of (13), the result follows from the law of iterated expectation. \square

Remark 4. Note that the previous subadditivity property does not hold in general. Lee and Prékopa (2013) provide a counterexample in the case where the underlying random vectors follow particular discrete distributions.

2.2. Archimedean copula case

In this section, we focus on multivariate random vectors distributed as particular Archimedean copulas. Note that, since Archimedean copulas are exchangeable, all components of the introduced multivariate risk measures are the same, when applied to random vectors with such distributions. We show that analytical expressions can be obtained for the lower-orthant and the upper-orthant CTE, which allows us to illustrate the behavior of the proposed risk-measures. In the next sections, the presented properties will be investigated more formerly.

We restrain ourself to random vectors with uniform marginals. In that case, the multivariate VaR-s introduced in Cousin and Di Bernardino (2013) can be expressed as an integral transformation of the Archimedean generator.

Definition 2.3. A d -dimensional Archimedean copula C with generator ϕ is a distribution function on $(0, 1)^d$ defined by

$$C(u_1, \dots, u_d) = \phi^{-1}(\phi(u_1) + \dots + \phi(u_d)),$$

where ϕ^{-1} is a d -monotone function from $[0, \infty)$ to $[0, 1]$.

Proposition 2.5. Let \mathbf{X} be a d -dimensional random vector which follows an Archimedean copula with generator ϕ , then, for any $i = 1, \dots, d$,

$$\underline{\text{VaR}}_\alpha^i(\mathbf{X}) = 1 - \int_\alpha^1 \left(1 - \frac{\phi(u)}{\phi(\alpha)}\right)^{d-1} du, \quad (14)$$

$$\overline{\text{VaR}}_\alpha^i(\mathbf{1} - \mathbf{X}) = \int_{1-\alpha}^1 \left(1 - \frac{\phi(u)}{\phi(1-\alpha)}\right)^{d-1} du. \quad (15)$$

Proof. This comes down from Corollary 8 in Cousin and Di Bernardino (2013). \square

Consequently, by using representations (8)–(9) and relations (14)–(15), the following integral expressions hold for any $i = 1, \dots, d$

$$\begin{aligned} \underline{\text{CTE}}_\alpha^i(\mathbf{X}) &= \frac{1}{1 - K(\alpha)} \\ &\times \int_\alpha^1 \left(1 - \int_\gamma^1 \left(1 - \frac{\phi(u)}{\phi(\gamma)}\right)^{d-1} du\right) \\ &\times K'(\gamma) d\gamma, \end{aligned} \quad (16)$$

$$\begin{aligned} \overline{\text{CTE}}_\alpha^i(\mathbf{1} - \mathbf{X}) &= \frac{1}{K(1-\alpha)} \\ &\times \int_\alpha^1 \left(\int_{1-\gamma}^1 \left(1 - \frac{\phi(u)}{\phi(1-\gamma)}\right)^{d-1} du\right) \\ &\times K'(1-\gamma) d\gamma, \end{aligned} \quad (17)$$

where K is the Kendall distribution function associated with X . Using (16), we are able to give analytical expressions of $\underline{\text{CTE}}_\alpha^i(\mathbf{X})$ for some particular families of Archimedean copulas. Additionally, from Example 1, analogous expressions can be obtained for the upper-orthant CTE of the vector $\mathbf{1} - \mathbf{X} = (1 - X_1, \dots, 1 - X_d)$ given that the Kendall distribution K of \mathbf{X} is known. One can use either formula (17) or the following relation

$$\begin{aligned} \overline{\text{CTE}}_\alpha^i(\mathbf{1} - \mathbf{X}) &= \frac{1}{K(1-\alpha)} \left[\frac{1}{2} - (1 - K(1-\alpha)) \right. \\ &\left. \times (1 - \underline{\text{CTE}}_{1-\alpha}^i(\mathbf{X})) \right] \end{aligned} \quad (18)$$

which comes down from Eq. (10). In Table 1, we provide analytical expressions of the Kendall distribution function $K(\alpha)$ for bivariate Gumbel, Frank, Clayton and Ali-Mikhail-Haq families.

As a matter of example, we now focus on Clayton and Gumbel copula families.

Clayton family illustration

Let us now consider the Clayton family of bivariate copulas. This family is interesting since it contains the counter-monotonic, the independence and the comonotonic copulas as particular cases. Clayton copulas are associated with generator ϕ of the form $\phi(u) = \frac{1}{\theta}(u^{-\theta} - 1)$, $u \in (0, 1)$ with a dependence parameter θ . Let (X, Y) be a random vector distributed as a Clayton copula with parameter $\theta \geq -1$. Then, X and Y are uniformly-distributed on $(0, 1)$ and the joint distribution function C_θ of (X, Y) is such that

$$\begin{aligned} C_\theta(x, y) &= (\max\{x^{-\theta} + y^{-\theta} - 1, 0\})^{-\frac{1}{\theta}}, \\ &\text{for } \theta \geq -1, (x, y) \in [0, 1]^2. \end{aligned} \quad (19)$$

We obtain in Table 2 a closed-form expression for the multivariate lower-orthant CTE in that case.

Interestingly, one can readily show that $\frac{\partial \underline{\text{CTE}}_{\alpha, \theta}^i}{\partial \alpha} \geq 0$ and $\frac{\partial \underline{\text{CTE}}_{\alpha, \theta}^i}{\partial \theta} \leq 0$, for $\theta \geq -1$ and $\alpha \in (0, 1)$. This proves that, for Clayton-distributed random vectors, the components of our CTE are increasing functions of the risk level α and decreasing functions of

Table 1
Kendall distribution in some classical bivariate Archimedean copulas.

Copula	$\theta \in$	Kendall distribution $K(\alpha, \theta)$
Gumbel	$[1, \infty)$	$\alpha \left(1 - \frac{1}{\theta} \ln \alpha\right)$
Frank	$(-\infty, \infty) \setminus \{0\}$	$\alpha + \frac{1}{\theta} (1 - e^{\theta\alpha}) \ln \left(\frac{1 - e^{-\theta\alpha}}{1 - e^{-\theta}}\right)$
Clayton	$[-1, \infty) \setminus \{0\}$	$\alpha \left(1 + \frac{1}{\theta} (1 - \alpha^\theta)\right)$
Ali–Mikhail–Haq	$[-1, 1)$	$\frac{\alpha - 1 + \theta + (1 - \theta + \theta\alpha)(\ln(1 - \theta + \theta\alpha) + \ln \alpha)}{\theta - 1}$

Table 2
 $\text{CTE}_{\alpha, \theta}^i(X, Y)$, $i = 1, 2$, for different copula dependence structures.

Copula	θ	$\text{CTE}_{\alpha, \theta}^i(X, Y)$
Clayton C_θ	$(-1, \infty)$	$\frac{1}{2} \frac{\theta - 1 - \alpha^2(1+\theta) + 2\alpha^{1+\theta}}{\theta - 1 - \alpha(1+\theta) + \alpha^{1+\theta}}$
Counter-monotonic W	-1	$\frac{1}{4} \frac{1 - \alpha^2 + 2 \ln \alpha}{1 - \alpha + \ln \alpha}$
Independent Π	0	$\frac{1}{2} \frac{(1 - \alpha)^2}{1 - \alpha + \alpha \ln \alpha}$
$\frac{\pi}{\Sigma - \Pi}$	1	$\frac{1}{2} \frac{1 + \alpha^2(2 \ln \alpha - 1)}{(1 - \alpha)^2}$
Comonotonic M	∞	$\frac{1 + \alpha}{2}$

the dependence parameter θ . Note also that, in the comonotonic case, our CTE corresponds to the vector composed of the univariate CTE associated with each component. These properties are illustrated in Fig. 1 (left panel) where $\text{CTE}_{\alpha, \theta}^i(X, Y)$ is plotted as a function of the risk level α for different values of θ . Observe that an increase of the dependence parameter θ tends to lower the CTE up to the perfect dependence case where $\text{CTE}_{\alpha, \theta = +\infty}^i(X, Y) = \text{CTE}_\alpha(X) = \frac{1 + \alpha}{2}$. The previous empirical behaviors will be formally investigated in next sections.

In the same framework, using Eq. (18), one can readily show that $\frac{\partial \text{CTE}_{\alpha, \theta}^i}{\partial \alpha} \geq 0$ and $\frac{\partial \text{CTE}_{\alpha, \theta}^i}{\partial \theta} \geq 0$, for $\theta \geq -1$ and $\alpha \in (0, 1)$. This proves that, for random couples with uniform margins and Clayton survival copula, the components of our multivariate CTE are increasing functions both of the risk level α and of the dependence parameter θ . Note also that the multivariate CTE in the comonotonic case corresponds to the vector composed of the univariate CTE associated with each component. These properties are illustrated in Fig. 1 (right panel) where $\overline{\text{CTE}}_{\alpha, \theta}^i(1 - X, 1 - Y)$ is plotted as a function of the risk level α for different values of the parameter θ . Observe that, contrary to the lower-orthant CTE , an increase of the dependence parameter θ tends to increase the $\overline{\text{CTE}}$. Then in the case of upper-orthant CTE , the upper bound is represented by the perfect dependence case where $\overline{\text{CTE}}_{\alpha, \theta = +\infty}^1(X, Y) = \text{CTE}_\alpha(X) = \frac{1 + \alpha}{2}$. The latter empirical behaviors will be formally confirmed in next sections.

Gumbel family illustration

Gumbel copulas are associated with a generator ϕ of the form $\phi(u) = (-\log(u))^\theta$, $u \in (0, 1)$ with a dependence parameter $\theta \geq 1$. The bivariate family of copulas is such that

$$C_\theta(x, y) = e^{-((-\ln x)^\theta + (-\ln y)^\theta)^{\frac{1}{\theta}}}$$

for $\theta \in [1, \infty)$, $(x, y) \in [0, 1]^2$ (e.g., see Section 3.3.1 in Nelsen, 1999) and X, Y standard uniform marginals. For $\theta = 1$ we have the independent copula $C_1(x, y) = \Pi(x, y) = xy$; for $\theta = \infty$ the Fréchet bound $M(x, y) = \min\{x, y\}$ (comonotonic random variables). Again, in this case, analytical expressions of $\text{CTE}_\alpha(X, Y)$ and $\overline{\text{CTE}}_{\alpha, \theta}^1(1 - X, 1 - Y)$ can be derived. As in the Clayton case, it can be

proved that $\frac{\partial \text{CTE}_{\alpha, \theta}^1}{\partial \alpha} \geq 0$, $\frac{\partial \text{CTE}_{\alpha, \theta}^1}{\partial \theta} \leq 0$, $\frac{\partial \overline{\text{CTE}}_{\alpha, \theta}^1}{\partial \alpha} \geq 0$ and $\frac{\partial \overline{\text{CTE}}_{\alpha, \theta}^1}{\partial \theta} \geq 0$, for $\theta \geq 1$ and $\alpha \in (0, 1)$. Fig. 2 plots one component of the proposed measures in the Gumbel case (all components are the same by exchangeability), as a function of the risk level α and for different dependence parameter θ . We can see that the behavior of the

upper-orthant and the lower-orthant CTE are similar to the one exhibited in Fig. 1 for the Clayton case.

2.3. Comparison with other risk measures

We now discuss whether the components of CTE and $\overline{\text{CTE}}$ can be compared with univariate VaR and CTE applied to the corresponding marginal risks and under which condition components of CTE and $\overline{\text{CTE}}$ are more conservative than the components of VaR and VaR.

Proposition 2.6. Consider a d -dimensional random vector \mathbf{X} . Assume that its multivariate distribution function F is quasi concave.⁶ Then, for any $i = 1, \dots, d$, the following inequality holds:

$$\text{VaR}_\alpha(X_i) \leq \text{CTE}_\alpha^i(\mathbf{X}), \quad \text{for } i = 1, \dots, d. \tag{20}$$

Proof. Let $\alpha \in (0, 1)$. From the definition of the accumulated probability, it is easy to show that $\partial L(\alpha)$ is inferiorly bounded by the marginal univariate quantile functions. Moreover, recall that $L(\alpha)$ is a convex set in \mathbb{R}_+^d from the quasi concavity of F (see Section 2 in Tibiletti, 1995). Then, for all $\mathbf{x} = (x_1, \dots, x_d) \in L(\alpha)$, $x_1 \geq \text{VaR}_\alpha(X_1), \dots, x_d \geq \text{VaR}_\alpha(X_d)$ and trivially, $\text{CTE}_\alpha^i(\mathbf{X})$ is greater than $\text{VaR}_\alpha(X_i)$, for $i = 1, \dots, d$. Hence the result. \square

Proposition 2.6 states that the multivariate lower-orthant CTE_α is more conservative than the vector composed of the classical univariate α -Value-at-Risk of marginal distributions. Such a result is not true for the upper-orthant CTE as can be seen in left panel of Figs. 1–2.

It is interesting to remark that, for comonotonic random vectors, the proposed multivariate CTE -s coincide with the vector composed of univariate CTE -s of the corresponding marginals. For the sake of clarity, we firstly recall the definition of comonotonicity for a random vector (see e.g., Definition 1.9.1 in Denuit et al., 2005).

Definition 2.4. A random vector $\mathbf{X} = (X_1, \dots, X_d)$ is comonotonic if and only if there exists a random variable Z and increasing functions g_1, \dots, g_d such that \mathbf{X} is equal to $(g_1(Z), \dots, g_d(Z))$ in distribution.

Proposition 2.7. Consider a comonotonic non-negative d -dimensional random vector \mathbf{X} . Then, for all $\alpha \in (0, 1)$, it holds that

$$\text{CTE}_\alpha^i(\mathbf{X}) = \text{CTE}_\alpha(X_i) = \overline{\text{CTE}}_\alpha^i(\mathbf{X}), \quad \text{for } i = 1, \dots, d.$$

Proof. From Definition 2.4, we write the distribution of vector \mathbf{X} as $F_{\mathbf{X}}(x) = F_Z(\min_{i=1, \dots, d}\{g_i^{-1}(x_i)\})$ and $\overline{F}_{\mathbf{X}}(x) = \overline{F}_Z(\max_{i=1, \dots, d}\{g_i^{-1}(x_i)\})$ so that $F_{\mathbf{X}}(\mathbf{X}) = F_Z(Z) = F_{X_i}(X_i)$ and $\overline{F}_{\mathbf{X}}(\mathbf{X}) = \overline{F}_Z(Z) = \overline{F}_{X_i}(X_i)$ for all $i = 1, \dots, d$. The result follows immediately. \square

Let us now compare the multivariate lower-orthant and upper-orthant CTE introduced in Definition 2.1 with the multivariate lower-orthant and upper-orthant VaR defined by Cousin and Di Bernardino (2013).

Proposition 2.8. Consider a d -dimensional random vector \mathbf{X} satisfying the regularity conditions. If $\text{VaR}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α , then it holds that

$$\text{CTE}_\alpha^i(\mathbf{X}) \geq \text{VaR}_\alpha^i(\mathbf{X}), \quad \text{for all } \alpha \in (0, 1).$$

⁶ A function F is quasi concave if for any x, y , $F(px + (1 - p)y) \geq \min(F(x), F(y))$ for all $p \in (0, 1)$. Note that a function F is quasi concave if and only if its upper level sets of are convex. Tibiletti (1995) points out families of distribution functions which satisfy the property of quasi concavity. For instance, multivariate elliptical distributions and Archimedean copulas are quasi concave functions.




Fig. 1. Behavior of $\underline{\text{CTE}}_{\alpha,\theta}^1(X, Y) = \underline{\text{CTE}}_{\alpha,\theta}^2(X, Y)$ (left) and $\overline{\text{CTE}}_{\alpha,\theta}^1(1-X, 1-Y) = \overline{\text{CTE}}_{\alpha,\theta}^2(1-X, 1-Y)$ (right) with respect to risk level α for different values of dependence parameter θ . The random vector (X, Y) follows a Clayton copula distribution with parameter θ . The horizontal line corresponds to $\mathbb{E}[X] = \frac{1}{2}$. Note that this limit is reached (right panel) for the upper-orthant CTE in the counter-monotonic case, i.e., $\overline{\text{CTE}}_{\alpha,\theta=-1}^1(1-X, 1-Y) = \frac{1}{2}$.




Fig. 2. Behavior of $\underline{\text{CTE}}_{\alpha,\theta}^1(X, Y) = \underline{\text{CTE}}_{\alpha,\theta}^2(X, Y)$ (left) and $\overline{\text{CTE}}_{\alpha,\theta}^1(1-X, 1-Y) = \overline{\text{CTE}}_{\alpha,\theta}^2(1-X, 1-Y)$ (right) with respect to risk level α for different values of dependence parameter θ . The random vector (X, Y) follows a Gumbel copula distribution with parameter θ . The horizontal line corresponds to $\mathbb{E}[X] = \frac{1}{2}$.

If $\overline{\text{VaR}}_{\alpha}^i(\mathbf{X})$ is a non-decreasing function of α , then it holds that

$$\overline{\text{CTE}}_{\alpha}^i(\mathbf{X}) \geq \overline{\text{VaR}}_{\alpha}^i(\mathbf{X}), \quad \text{for all } \alpha \in (0, 1).$$

Proof. From Eq. (8), since $\overline{\text{VaR}}_{\gamma}^i(\mathbf{X})$ is assumed to be non-decreasing with respect to γ , we obtain

$$\begin{aligned} \overline{\text{CTE}}_{\alpha}^i(\mathbf{X}) &\geq \frac{1}{1-K(\alpha)} \int_{\alpha}^1 \overline{\text{VaR}}_{\gamma}^i(\mathbf{X}) K'(\gamma) d\gamma \\ &= \frac{K(1) - K(\alpha)}{1 - K(\alpha)} \overline{\text{VaR}}_{\alpha}^i(\mathbf{X}) = \overline{\text{VaR}}_{\alpha}^i(\mathbf{X}). \end{aligned}$$

From Eq. (9), if $\overline{\text{VaR}}_{\gamma}^i(\mathbf{X})$ is assumed to be non-decreasing with respect to γ , we obtain

$$\begin{aligned} \overline{\text{CTE}}_{\alpha}^i(\mathbf{X}) &\geq \frac{1}{\widehat{K}(1-\alpha)} \int_{\alpha}^1 \overline{\text{VaR}}_{\gamma}^i(\mathbf{X}) \widehat{K}'(1-\gamma) d\gamma \\ &= \frac{\widehat{K}(1-\alpha) - \widehat{K}(0)}{\widehat{K}(1-\alpha)} \overline{\text{VaR}}_{\alpha}^i(\mathbf{X}) = \overline{\text{VaR}}_{\alpha}^i(\mathbf{X}). \end{aligned}$$

Hence the result. \square

This property holds for instance for any random vector which admits an Archimedean copula. It is indeed proved in Cousin and Di

Bernardino (2013) that $\text{VaR}_{\alpha}(\mathbf{X})$ and $\overline{\text{VaR}}_{\alpha}(\mathbf{X})$ are non-decreasing functions of the risk level α for such dependence structures.

2.4. Behavior of multivariate CTE with respect to marginal distributions

In this section we study the behavior of the multivariate lower-orthant and upper-orthant Conditional-Tail-Expectation with respect to a change in marginal distributions. Results presented below provide a natural multivariate extension of classical results in a univariate setting (see, e.g., Denuit and Charpentier, 2004).

Proposition 2.9. Let \mathbf{X} and \mathbf{Y} be two d -dimensional random vector satisfying the regularity conditions and with the same copula structure C . If $X_i \stackrel{d}{=} Y_i$, then it holds that

$$\underline{\text{CTE}}_{\alpha}^i(\mathbf{X}) = \underline{\text{CTE}}_{\alpha}^i(\mathbf{Y}), \quad \text{for all } \alpha \in (0, 1),$$

and

$$\overline{\text{CTE}}_{\alpha}^i(\mathbf{X}) = \overline{\text{CTE}}_{\alpha}^i(\mathbf{Y}), \quad \text{for all } \alpha \in (0, 1).$$

Proof. From Proposition 2.6 in Cousin and Di Bernardino (2013), we know that $\text{VaR}_{\alpha}^i(\mathbf{X}) = \text{VaR}_{\alpha}^i(\mathbf{Y})$ and $\overline{\text{VaR}}_{\alpha}^i(\mathbf{X}) = \overline{\text{VaR}}_{\alpha}^i(\mathbf{Y})$. The demonstration comes down from representation (8) and (9). \square

Table 3

$\text{CTE}_\alpha(X, Y_i)$, for $i = 1, \dots, 5$, with the same copula Clayton copula with parameter 1, $X \sim \text{Exp}(1)$ and $Y_1 \sim \text{Exp}(2)$; $Y_2 \sim \text{Burr}(2, 1)$; $Y_3 \sim \text{Exp}(1)$; $Y_4 \sim \text{Fréchet}(4)$; $Y_5 \sim \text{Burr}(4, 1)$. This table provides an illustration of Propositions 2.9–2.10.

α	$\text{CTE}_\alpha(X, Y_1)$	$\text{CTE}_\alpha(X, Y_2)$	$\text{CTE}_\alpha(X, Y_3)$	$\text{CTE}_\alpha(X, Y_4)$	$\text{CTE}_\alpha(X, Y_5)$
0.10	(1.188, 0.594)	(1.188, 1.838)	(1.188, 1.188)	(1.188, 1.315)	(1.188, 1.229)
0.24	(1.449, 0.724)	(1.449, 2.218)	(1.449, 1.449)	(1.449, 1.431)	(1.449, 1.366)
0.38	(1.727, 0.864)	(1.727, 2.661)	(1.727, 1.727)	(1.727, 1.555)	(1.727, 1.506)
0.52	(2.049, 1.025)	(2.049, 3.235)	(2.049, 2.049)	(2.049, 1.704)	(2.049, 1.667)
0.66	(2.454, 1.227)	(2.454, 4.074)	(2.454, 2.454)	(2.454, 1.902)	(2.454, 1.876)
0.80	(3.039, 1.519)	(3.039, 5.591)	(3.039, 3.039)	(3.039, 2.219)	(3.039, 2.202)
0.90	(3.768, 1.884)	(3.768, 8.175)	(3.768, 3.768)	(3.768, 2.675)	(3.768, 2.665)
0.99	(6.102, 3.059)	(6.102, 26.59)	(6.102, 6.102)	(6.102, 4.813)	(6.102, 4.811)

In particular, when one component of the underlying risk vector changes, it does not affect the other components of the multivariate CTE-s as far as the dependence structure is unchanged. In the following, we analyze how our multivariate measures behave when the marginal risks increase with respect to the first-order stochastic dominance (e.g., see Definition 3.3.1. in Denuit et al., 2005). We recall below the definition of the first-order stochastic dominance.

Definition 2.5 (First-Order Stochastic Dominance). Let X and Y be two random variables. Then X is said to be smaller than Y in first-order stochastic dominance, denoted as $X \leq_{st} Y$, if the inequality

$$Q_X(\alpha) \leq Q_Y(\alpha) \text{ is satisfied for all } \alpha \in (0, 1),$$

where $Q_X(\alpha)$ (resp. $Q_Y(\alpha)$) is the α -th quantile of the distribution of X (resp. of Y).

Proposition 2.10. Let \mathbf{X} and \mathbf{Y} be two d -dimensional random vectors satisfying the regularity conditions and with the same copula structure C . If $X_i \leq_{st} Y_i$ as in Definition 2.5, then it holds that

$$\text{CTE}_\alpha^i(\mathbf{X}) \leq \text{CTE}_\alpha^i(\mathbf{Y}), \text{ for all } \alpha \in (0, 1),$$

and

$$\overline{\text{CTE}}_\alpha^i(\mathbf{X}) \leq \overline{\text{CTE}}_\alpha^i(\mathbf{Y}), \text{ for all } \alpha \in (0, 1).$$

Proof. From Proposition 2.7 in Cousin and Di Bernardino (2013), we know that, if $X_i \leq_{st} Y_i$ then $\text{VaR}_\alpha^i(\mathbf{X}) \leq \text{VaR}_\alpha^i(\mathbf{Y})$ and $\overline{\text{VaR}}_\alpha^i(\mathbf{X}) \leq \overline{\text{VaR}}_\alpha^i(\mathbf{Y})$. The result comes down from representation (8) and (9). \square

Example 2. In this example, we provide an illustration of Propositions 2.9–2.10 above. We consider the case of lower-orthant Conditional-Tail-Expectation (the upper-orthant case is completely analogous). The obtained results are gathered in Table 3.

We consider five different bivariate random vectors (X, Y_i) , for $i = 1, \dots, 5$, with the same bivariate Clayton copula with parameter 1. Let

$$\begin{aligned} X &\sim \text{Exp}(1), & Y_1 &\sim \text{Exp}(2), & Y_2 &\sim \text{Burr}(2, 1), \\ Y_3 &\sim \text{Exp}(1), & Y_4 &\sim \text{Fréchet}(4) & \text{and } Y_5 &\sim \text{Burr}(4, 1). \end{aligned}$$

We calculate $\text{CTE}_\alpha(X, Y_i)$, for $i = 1, \dots, 5$ (see Table 3). As proved in Proposition 2.9 we obtain an invariant property on the first coordinate of all $\text{CTE}_\alpha(X, Y_i)$, for $i = 1, \dots, 5$. Furthermore, since (X, Y_3) is an exchangeable continuous random vector, then $\text{CTE}_\alpha^1(X, Y_3) = \text{CTE}_\alpha^2(X, Y_3)$. Moreover, as $Q_X(\alpha) = 2Q_{Y_1}(\alpha)$, then $\text{CTE}_\alpha^1(X, Y_1) = 2\text{CTE}_\alpha^2(X, Y_1)$, for $\alpha \in (0, 1)$ (see Table 3).

By in Proposition 2.10, since $Y_1 \leq_{st} Y_5 \leq_{st} Y_4 \leq_{st} Y_2$, then,

$$\begin{aligned} \text{CTE}_\alpha^2(X, Y_1) &\leq \text{CTE}_\alpha^2(X, Y_5) \leq \text{CTE}_\alpha^2(X, Y_4) \\ &\leq \text{CTE}_\alpha^2(X, Y_2), \text{ for any level } \alpha \in (0, 1). \end{aligned}$$

Analogously, we also obtain $\text{CTE}_\alpha(X, Y_1) \leq \text{CTE}_\alpha(X, Y_3) \leq \text{CTE}_\alpha(X, Y_2)$, for any level $\alpha \in (0, 1)$.

Conversely Y_3, Y_4 and Y_3, Y_5 are not ordered in stochastic dominance sense.

In the following proposition, we investigate the effect of a mean-preserving change in marginal risks.

Proposition 2.11. Let \mathbf{X} and \mathbf{Y} be two d -dimensional random vectors satisfying the regularity conditions and with the same copula structure C . Let us assume that $\mathbb{E}[X_i] \leq \mathbb{E}[Y_i]$ and that the i th components of the lower-orthant VaR of X and Y satisfy the following single-cut condition: there exists a real c in $[0, 1)$ such that $\text{VaR}_\alpha^i(\mathbf{X}) \geq \text{VaR}_\alpha^i(\mathbf{Y})$ for all $\alpha < c$ and $\text{VaR}_\alpha^i(\mathbf{X}) \leq \text{VaR}_\alpha^i(\mathbf{Y})$ for all $\alpha \geq c$. Then,

$$\text{CTE}_\alpha^i(\mathbf{X}) \leq \text{CTE}_\alpha^i(\mathbf{Y}), \text{ for all } \alpha \in (0, 1).$$

The same result holds $\overline{\text{CTE}}$.

Proof. Let us consider the function Δ defined by

$$\begin{aligned} \alpha \mapsto \Delta(\alpha) &= (1 - K(\alpha)) (\text{CTE}_\alpha^i(\mathbf{Y}) - \text{CTE}_\alpha^i(\mathbf{X})) \\ &= \int_\alpha^1 (\text{VaR}_\gamma^i(\mathbf{Y}) - \text{VaR}_\gamma^i(\mathbf{X})) K'(\gamma) d\gamma. \end{aligned}$$

Then $\Delta(0) = (1 - K(0))(\mathbb{E}[Y_i] - \mathbb{E}[X_i]) \geq 0$, $\Delta(1) = 0$ and $\Delta'(\alpha) = (\text{VaR}_\alpha^i(\mathbf{X}) - \text{VaR}_\alpha^i(\mathbf{Y}))K'(\alpha)$ is positive for all $\alpha < c$ and negative for all $\alpha \geq c$ thanks to the single-cut condition. As a result, the function Δ remains positive on $[0, 1]$ and the result holds. \square

This result is similar to the consistency property of univariate CTE with respect to the less dangerous order, i.e., for univariate random variables X and Y such as $X \leq_D Y$, then, for any α in $[0, 1]$, $\text{CTE}_\alpha(X) \leq \text{CTE}_\alpha(Y)$.

Example 3. We provide here an illustration of Proposition 2.11. Firstly we remark in Example 2 that $\mathbb{E}[Y_3] < \mathbb{E}[Y_5]$. However, the single-cut condition of Proposition 2.11 for $\text{VaR}_\alpha^2(X, Y_3)$ and $\text{VaR}_\alpha^2(X, Y_5)$ is not satisfied and from Table 3 one can see that $\text{CTE}_\alpha^2(X, Y_3)$, $\text{CTE}_\alpha^2(X, Y_5)$ are not ordered for any level $\alpha \in (0, 1)$.

We now define $\mathbf{X} := (X, Z_1)$ and $\mathbf{Y} := (X, \frac{1}{2}(Z_1 + Z_2))$. Let X a continuous random variable. Let Z_1, Z_2 be two independent random variables such that $Z_1 \sim Z_2 \sim \text{Exp}(2)$, then $Z_1 + Z_2 \sim \text{Erlang}(2, 2)$ distribution. Assume that \mathbf{X} and \mathbf{Y} have both the same independent copula C . One can show that $\mathbb{E}[\frac{1}{2}(Z_1 + Z_2)] = \mathbb{E}[Z_1] = \frac{1}{2}$ and there exists a real c in $[0, 1)$ such that $\text{VaR}_\alpha^2(\mathbf{Y}) \geq \text{VaR}_\alpha^2(\mathbf{X})$ for all $\alpha < c$ and $\text{VaR}_\alpha^2(\mathbf{Y}) \leq \text{VaR}_\alpha^2(\mathbf{X})$ for all $\alpha \geq c$ (see Fig. 3, left). The interested reader is also referred to Section 7.3 in Denuit et al. (2005).

Then, using Proposition 2.11 we obtain $\text{CTE}_\alpha^2(\mathbf{Y}) \leq \text{CTE}_\alpha^2(\mathbf{X})$, for all $\alpha \in (0, 1)$. This property is illustrated in Fig. 3 (right).

2.5. Behavior of multivariate CTE with respect to the dependence structure

In this section we study the behavior of our CTE generalizations with respect to a variation of the dependence structure, with unchanged marginal distributions.




Fig. 3. Left: $\text{VaR}_\alpha^2(\mathbf{X})$ (full line) and $\text{VaR}_\alpha^2(\mathbf{Y})$ (dashed line). We also represent the associated univariate VaR, i.e. $\text{VaR}_\alpha(Z_1)$ (squared line) and $\text{VaR}_\alpha(\frac{1}{2}(Z_1 + Z_2))$ (dotted line). Remark that $\text{VaR}_\alpha^2(\mathbf{X})$ and $\text{VaR}_\alpha^2(\mathbf{Y})$ verify the single-cut condition in Proposition 2.11. Right: $\text{CTE}_\alpha^2(\mathbf{X})$ (full line) and $\text{CTE}_\alpha^2(\mathbf{Y})$ (dashed line).

Proposition 2.12. Let \mathbf{X} and \mathbf{X}^* be two d -dimensional continuous random vectors satisfying the regularity conditions and with the same margins F_{X_i} and $F_{X_i^*}$, for $i = 1, \dots, d$, and let C (resp. C^*) denote the copula function associated with \mathbf{X} (resp. \mathbf{X}^*) and \bar{C} (resp. \bar{C}^*) the survival copula function associated with \mathbf{X} (resp. \mathbf{X}^*).

Let $U_i = F_{X_i}(X_i)$, $U_i^* = F_{X_i^*}(X_i^*)$, $\mathbf{U} = (U_1, \dots, U_d)$ and $\mathbf{U}^* = (U_1^*, \dots, U_d^*)$.

If $[U_i | C(\mathbf{U}) \geq \alpha] \leq_{\text{st}} [U_i^* | C^*(\mathbf{U}^*) \geq \alpha]$
 then $\text{CTE}_\alpha^i(\mathbf{X}) \leq \text{CTE}_\alpha^i(\mathbf{X}^*)$.

Let $V_i = \bar{F}_{X_i}(X_i)$, $V_i^* = \bar{F}_{X_i^*}(X_i^*)$, $\mathbf{V} = (V_1, \dots, V_d)$ and $\mathbf{V}^* = (V_1^*, \dots, V_d^*)$.

If $[V_i | \bar{C}(\mathbf{V}) \leq 1 - \alpha] \leq_{\text{st}} [V_i^* | \bar{C}^*(\mathbf{V}^*) \leq 1 - \alpha]$
 then $\overline{\text{CTE}}_\alpha^i(\mathbf{X}) \geq \overline{\text{CTE}}_\alpha^i(\mathbf{X}^*)$.

Proof. We recall that $U_1 \leq_{\text{st}} U_2$ if and only if $\mathbb{E}[f(U_1)] \leq \mathbb{E}[f(U_2)]$, for all non-decreasing function f , such that the expectations exist (see Denuit et al., 2005, Proposition 3.3.14). We now choose $f(u) = Q_{X_i}(u)$, for $u \in (0, 1)$. Then we obtain

$$\mathbb{E}[Q_{X_i}(U_i) | C(\mathbf{U}) \geq \alpha] \leq \mathbb{E}[Q_{X_i}(U_i^*) | C^*(\mathbf{U}^*) \geq \alpha].$$

But the right-hand side of the previous inequality is equal to $\mathbb{E}[Q_{X_i^*}(U_i^*) | C^*(\mathbf{U}^*) \geq \alpha]$ since X_i and X_i^* have the same distribution. Finally, we obtain $\text{CTE}_\alpha^i(\mathbf{X}) \leq \text{CTE}_\alpha^i(\mathbf{X}^*)$. For the second point of the statement we choose the non-decreasing function $f(u) = -\bar{F}_{X_i}^{-1}(u)$, for $u \in (0, 1)$. Since X_i and X_i^* have the same distribution, we obtain the result. \square

We now provide an illustration of Proposition 2.12 in the case of bivariate Archimedean copulas.

Corollary 2.2. Consider a 2-dimensional random vector \mathbf{X} , satisfying the regularity conditions, with marginal distributions F_{X_i} , for $i = 1, \dots, d$, copula C and survival copula \bar{C} .

If C belongs to one of the bivariate family of Archimedean copulas introduced in Table 1, an increase of the dependence parameter θ yields a decrease in each component of $\text{CTE}_\alpha(\mathbf{X})$.

If \bar{C} belongs to one of the bivariate family of Archimedean copulas introduced in Table 1, an increase of the dependence parameter θ yields an increase in each component of $\overline{\text{CTE}}_\alpha(\mathbf{X})$.

Proof. In the bivariate Archimedean case, the joint distribution of $(U, C(U, V))$ can be obtained analytically by using a change of variable transformation⁷ from (U, V) to $(U, C(U, V))$:

$$F_{(U, C(U, V))}(u, \alpha) = \alpha - \frac{\phi(\alpha)}{\phi'(\alpha)} + \frac{\phi(u)}{\phi'(\alpha)}, \quad 0 < \alpha < u < 1. \quad (21)$$

Then, thanks to formula (21), we can obtain the joint survival probability

$$\begin{aligned} h(u, \alpha) &:= \mathbb{P}[U \geq u, C(U, V) \geq \alpha] \\ &= 1 - u + \frac{\phi(u)}{\phi'(\alpha)} \quad \text{for } 0 < \alpha < u < 1. \end{aligned} \quad (22)$$

Note that $1 - K(\alpha) = \mathbb{P}[C(U, V) \geq \alpha] = h(\alpha, \alpha)$. Let C_θ and C_{θ^*} be two bivariate Archimedean copulas of the same family with generator ϕ_θ and ϕ_{θ^*} such that $\theta \leq \theta^*$. Given Proposition 2.12 and by exchangeability, we only have to check that the relation $[U^* | C_{\theta^*}(U^*, V^*) \geq \alpha] \leq_{\text{st}} [U | C_\theta(U, V) \geq \alpha]$ hold where (U, V) and (U^*, V^*) are distributed according to (resp.) C_θ and C_{θ^*} . Given formula (22), the previous relation can be restated as

$$\frac{h^*(u, \alpha)}{h^*(\alpha, \alpha)} \leq \frac{h(u, \alpha)}{h(\alpha, \alpha)}, \quad \text{for } 0 < \alpha < u < 1, \quad (23)$$

where, from (22), $h(u, \alpha) = 1 - u + \phi_\theta(u)/\phi'_\theta(\alpha)$ and $h^*(u, \alpha) = 1 - u + \phi_{\theta^*}(u)/\phi'_{\theta^*}(\alpha)$. Eventually, we have checked that, for all Archimedean family introduced in Table 1, Eq. (23) is indeed satisfied when $\theta \leq \theta^*$. We then immediately obtain from Proposition 2.12 that each component of $\text{CTE}_\alpha(\mathbf{X})$ is a decreasing function of θ . Analogously one can prove the result for $\overline{\text{CTE}}_\alpha(\mathbf{X})$. \square

Then, for copulas in Table 1, the multivariate lower-orthant CTE (resp. the upper-orthant $\overline{\text{CTE}}$) is non-increasing (non-decreasing) with respect to the dependence parameter θ .

In particular, this means that limit behaviors of dependence parameters are associated with bounds for CTE and $\overline{\text{CTE}}$ within the same family of Archimedean copulas.

2.6. Behavior of multivariate CTE with respect to risk level

We now study the behavior of the multivariate lower-orthant and upper-orthant Conditional-Tail-Expectation with respect to risk-level α . In particular we obtain the following result.

⁷ In the book by Nelsen (1999, Corollary 4.3.5), a geometrical argument is used instead to obtain the distribution function of $(U, C(U, V))$.

Corollary 2.3. Consider a d -dimensional random vector \mathbf{X} satisfying the regularity conditions.

1. If $\text{VaR}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α , then $\underline{\text{CTE}}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α .
2. If $\overline{\text{VaR}}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α , then $\overline{\text{CTE}}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α .

Proof. Let us consider the i th coordinate $\underline{\text{CTE}}_\alpha^i(\mathbf{X})$. From (8) we have

$$\frac{d}{d\alpha} \underline{\text{CTE}}_\alpha^i(\mathbf{X}) = \frac{K'(\alpha)}{1 - K(\alpha)} [\underline{\text{CTE}}_\alpha^i(\mathbf{X}) - \text{VaR}_\alpha^i(\mathbf{X})].$$

Using Proposition 2.8 the latter expression is non-negative for any level $\alpha \in (0, 1)$. The second point of Corollary 2.3 comes down analogously. \square

The following result proves that assumptions of Corollary 2.3 are satisfied in the large class of d -dimensional Archimedean copulas.

Corollary 2.4. Consider a d -dimensional random vector \mathbf{X} , satisfying the regularity conditions with copula C and survival copula \bar{C} .

1. If C is a d -dimensional Archimedean copula, then $\underline{\text{CTE}}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α .
2. If \bar{C} is a d -dimensional Archimedean copula, then $\overline{\text{CTE}}_\alpha^i(\mathbf{X})$ is a non-decreasing function of α .

Proof. The demonstration of this result comes down from Corollary 2.5 in Cousin and Di Bernardino (2013). \square

In the univariate setting, the Conditional-Tail-Expectation contains a safety loading i.e., $\text{CTE}_\alpha(X) \geq \mathbb{E}[X]$, $\forall \alpha \in (0, 1)$ (see Section 2.4.3.3 in Denuit et al., 2005). The safety loading should cover the fluctuations of loss experience. Corollary 2.5 below provides a similar property also for our multivariate lower-orthant and upper-orthant CTE.

Corollary 2.5. Under assumptions of Corollary 2.3, it holds that

$$\underline{\text{CTE}}_\alpha^i(\mathbf{X}) \geq \mathbb{E}[X_i], \quad \overline{\text{CTE}}_\alpha^i(\mathbf{X}) \geq \mathbb{E}[X_i],$$

for all $\alpha \in (0, 1)$.

Conclusion and perspectives

In this paper, we provide two extensions of the Conditional-Tail-Expectation to a multivariate setting. The lower-orthant CTE and the upper-orthant CTE can be viewed as natural counterparts of the lower-orthant VaR and upper-orthant VaR introduced in Cousin and Di Bernardino (2013). These measures transform risk vectors into real-valued vectors with the same dimension. The proposed multivariate CTE measures incorporate the entire extreme quadrant part of the underlying distribution contrary to their multivariate VaR counterparts. They are well-suited for intrinsically multivariate risk problems where, for instance, risks cannot be aggregated together or even compared. We have shown that most properties satisfy by the aforementioned multivariate VaR-s also hold for the studied multivariate CTE-s. In particular, the proposed CTE-s satisfy multivariate version of the positive homogeneity and the translation invariance property, which is not the case for instance for the classical Euler allocation measures recalled in the introduction of this paper. We show that these measures are additive for π -comonotonic couple of random vectors and we provide an evidence of the sub-additivity property in the independent case. Unsurprisingly, the behavior of the lower-orthant CTE (resp. upper-orthant CTE) with respect to a change in marginal risks or to

a change in dependence structure tends to be the same as for the lower-orthant VaR (resp. upper-orthant VaR).

Extension of discrete distributions can be done by adapting the work by Lee and Prékopa (2013). However this interesting topic goes beyond the scope of the present paper. Another subject of future research should be to compare our multivariate Conditional-Tail-Expectations with existing multivariate generalizations of these measures presented in the introduction, both theoretically and experimentally.

Acknowledgments

The authors thank an anonymous referee for constructive remarks to improve the paper. This work has been partially supported by the French research national agency (ANR) under the reference ANR-08BLAN-0314-01. Part of this work also benefit from the support of the MIRACCLE-GICC project.

References

- Acharya, V.V., Pedersen, L.H., Philippon, T., Richardson, M.P., 2010. Measuring systemic risk. FRB of Cleveland Working Paper.
- Adrian, T., Brunnermeier, M.K., 2011. CoVaR. NBER Working Paper.
- Artzner, P., Delbaen, F., Eber, J.M., Heath, D., 1999. Coherent measures of risk. *Math. Finance* 9 (3), 203–228.
- Bargès, M., Cossette, H., Marceau, É., 2009. TVaR-based capital allocation with copulas. *Insurance Math. Econom.* 45 (3), 348–361.
- Brownlees, C.T., Engle, R.F., 2012. Volatility, correlation and tails for systemic risk measurement. Working Paper.
- Burgert, C., Rüschendorf, L., 2006. Consistent risk measures for portfolio vectors. *Insurance Math. Econom.* 38 (2), 289–297.
- Cai, J.-J., Einmahl, J.H.J., de Haan, L., Zhou, C., 2013. Estimation of the marginal expected shortfall: the mean when a related variable is extreme. *CentER Discussion Paper*, 2012-080.
- Cai, J., Li, H., 2005. Conditional tail expectations for multivariate phase-type distributions. *J. Appl. Probab.* 42 (3), 810–825.
- Casco, I., Molchanov, I., 2007. Multivariate risks and depth-trimmed regions. *Finance Stoch.* 11 (3), 373–397.
- Chakak, A., Ezzer, M., 2000. Bivariate contours of copula. *Comm. Statist. Simulation Comput.* 29 (1), 175–185.
- Chaouch, M., Gannoun, A., Saracco, J., 2009. Estimation de quantiles géométriques conditionnels et non conditionnels. *J. SFDs* 150 (2), 1–27.
- Chauvigny, P., Devineau, L., Loisel, L., Maume-Deschamps, V., 2011. Fast remote but not extreme quantiles with multiple factors: applications to solvency ii and enterprise risk management. *Eur. Actuar. J.* 1 (1), 131–157.
- Cousin, A., Di Bernardino, E., 2013. On multivariate extensions of value-at-risk. *J. Multivariate Anal.* 119, 32–46.
- Denuit, M., Charpentier, A., 2004. *Mathématiques de l'Assurance non-vie. Tome 1: Principes Fondamentaux de Théorie du Risque*. Economica.
- Denuit, M., Dhaene, J., Goovaerts, M., Kaas, R., 2005. *Actuarial Theory for Dependent Risks*. Wiley.
- Di Bernardino, E., Laloë, T., Maume-Deschamps, V., Prieur, C., 2013. Plug-in estimation of level sets in a non-compact setting with applications in multivariate risk theory. *ESAIM Probab. Stat.* 17, 236–256.
- Ekeland, I., Galichon, A., Henry, M., 2012. Comonotonic measures of multivariate risks. *Math. Finance* 22 (1), 109–132.
- Embrechts, P., Puccetti, G., 2006. Bounds for functions of multivariate risks. *J. Multivariate Anal.* 97 (2), 526–547.
- Hamel, A.H., Heyde, F., 2010. Duality for set-valued measures of risk. *SIAM J. Financial Math.* 1 (1), 66–95.
- Imlahi, L., Ezzer, M., Chakak, A., 1999. Estimación de la curva mediana de una cópula $C(x_1, \dots, x_n)$. *Rev. R. Acad. Cienc. Exactas Fis. Nat.* 93 (2), 241–250.
- Jouini, E., Meddeb, M., Touzi, N., 2004. Vector-valued coherent risk measures. *Finance Stoch.* 8 (4), 531–552.
- Koltchinskii, V.I., 1997. M -estimation, convexity and quantiles. *Ann. Statist.* 25 (2), 435–477.
- Landsman, Z.M., Valdez, E.A., 2003. Tail conditional expectations for elliptical distributions. *N. Am. Actuar. J.* 7 (4), 55–71.
- Lee, J., Prékopa, A., 2013. Properties and calculation of multivariate risk measures: MVaR and MCVaR. *Ann. Oper. Res.* 211 (1), 225–254.
- Mainik, G., Schaanning, E., 2012. On dependence consistency of CoVaR and some other systemic risk measures. Working Paper.
- Massé, J.-C., Theodorescu, R., 1994. Halfplane trimming for bivariate distributions. *J. Multivariate Anal.* 48 (2), 188–202.
- Nappo, G., Spizzichino, F., 2009. Kendall distributions and level sets in bivariate exchangeable survival models. *Inform. Sci.* 179, 2878–2890.
- Nelsen, R.B., 1999. *An Introduction to Copulas*. In: *Lecture Notes in Statistics*, vol. 139. Springer-Verlag, New York.
- Prékopa, A., 2012. Multivariate value at risk and related topics. *Ann. Oper. Res.* 193, 49–69.
- Puccetti, G., Scarsini, M., 2010. Multivariate comonotonicity. *J. Multivariate Anal.* 101, 291–304.
- Rossi, C., 1973. Sulle curve di livello di una superficie di ripartizione in due variabili. *G. Ist. Ital. Degli Attuari* 36, 87–108.

- Rüschendorf, L., 2006. Law invariant convex risk measures for portfolio vectors. *Statist. Decisions* 24, 97–108.
- Scaillet, O., 2004. Nonparametric estimation and sensibility analysis of expected shortfall. *Math. Finance* 14 (1), 115–129.
- Serfling, R., 2002. Quantile functions for multivariate analysis: approaches and applications. *Statist. Neerlandica* 56 (2), 214–232. Special issue: Frontier research in theoretical statistics, 2000 (Eindhoven).
- Tasche, D., 2008. Capital allocation to business units and sub-portfolios: the Euler principle. *Quantitative finance Papers*, arXiv.org.
- Tibiletti, L., 1991. Sulla quasi concavità delle funzioni di ripartizione n -dimensionali—on quasi-concavity of n -dimensional distribution functions. In: *Atti del XV Convegno A.M.A.S.E.S.*, pp. 503–515.
- Tibiletti, L., 1995. Quasi-concavity property of multivariate distribution functions. *Ratio Math.* 9, 27–36.